



Official reprint from UpToDate®

[www.uptodate.com](http://www.uptodate.com) ©2021 UpToDate, Inc. and/or its affiliates. All Rights Reserved.

The content on the UpToDate website is not intended nor recommended as a substitute for medical advice, diagnosis, or treatment. Always seek the advice of your own physician or other qualified health care professional regarding any medical questions or conditions. The use of UpToDate content is governed by the [UpToDate Terms of Use](#). ©2021 UpToDate, Inc. All rights reserved.

## Clopidogrel: Patient drug information

Copyright 1978-2021 Lexicomp, Inc. All rights reserved.

(For additional information [see "Clopidogrel: Drug information"](#) and [see "Clopidogrel: Pediatric drug information"](#))

### Brand Names: US

Plavix

### Brand Names: Canada

ACT Clopidogrel; APO-Clopidogrel; Auro-Clopidogrel; BIO-Clopidogrel; DOM-Clopidogrel; JAMP-Clopidogrel; M-Clopidogrel; Mar-Clopidogrel; MINT-Clopidogrel [DSC]; MYLAN-Clopidogrel [DSC]; NRA-Clopidogrel; Plavix; PMS-Clopidogrel; PRIVA-Clopidogrel; RIVA-Clopidogrel; SANDOZ Clopidogrel; TARO-Clopidogrel; TEVA-Clopidogrel

### Warning

- This drug may not work as well in some people. A test can be done to see if you are one of these people. Talk with your doctor.

### What is this drug used for?

- It is used to lower the chance of heart attack or stroke.
- It may be given to you for other reasons. Talk with the doctor.

### What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have any of these health problems: Active bleeding or bleeding problems like bleeding in the brain or bleeding ulcers.
- If you are taking any of these drugs: Esomeprazole, omeprazole, or repaglinide.
- If you are taking any drugs that may raise the chance of bleeding. There are many drugs that can do this. Ask your doctor or pharmacist if you are not sure.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

## What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. This drug may need to be stopped before certain types of surgery as your doctor has told you. If this drug is stopped, your doctor will tell you when to start taking this drug again after your surgery or procedure.
- Do not stop taking this drug without calling the doctor who ordered it for you.
- If you fall or hurt yourself, or if you hit your head, call your doctor right away. Talk with your doctor even if you feel fine.
- You may bleed more easily. Be careful and avoid injury. Use a soft toothbrush and an electric razor. Rarely, some bleeding problems have been deadly.
- Talk with your doctor before using products that have aspirin, blood thinners, garlic, ginseng, ginkgo, ibuprofen or like products, pain drugs, or vitamin E.
- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

## What are some side effects that I need to call my doctor about right away?

**WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of bleeding like throwing up or coughing up blood; vomit that looks like coffee grounds; blood in the urine; black, red, or tarry stools; bleeding from the gums; abnormal vaginal bleeding; bruises without a cause or that get bigger; or bleeding you cannot stop.
- Signs of low blood sugar like dizziness, headache, feeling sleepy, feeling weak, shaking, a fast heartbeat, confusion, hunger, or sweating.
- A severe blood problem called thrombotic thrombocytopenic purpura (TTP) has happened with this drug. Sometimes, this has been deadly. Call your doctor right away if you feel confused, very tired, or very weak, or if you have weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or change in eyesight. Call your doctor right away if you have a fast heartbeat; fever; headache; pale skin; less urine passed or urine that is pink or has blood in it; purple spots on the skin or mouth; seizures; severe diarrhea, stomach pain, upset stomach, or throwing up; shortness of breath; or yellow skin or eyes.

## What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if you have any side effects that bother you or do not go away.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

## How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Take with or without food.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.

## What do I do if I miss a dose?

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses unless told to do so by your doctor.

## How do I store and/or throw out this drug?

- Store at room temperature in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

## General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Use of UpToDate is subject to the [Subscription and License Agreement](#).

Topic 12311 Version 156.0

→